arXiv:1105.3070v1 [math.GT] 16 May 2011

TEICHMÜLLER SPACE HAS NON-BUSEMANN POINTS

HIDEKI MIYACHI

Abstract. In this paper, we shall show that the metric boundary of the Teichmüller space with respect to the Teichmüller distance contains non-Busemann points when the complex dimension of the Teichmüller space is at least two.

1. Introduction

1.1. **Metric boundary and horofunction boundary.** The *metric boundary* of a metric space was defined by M. Rieffel in [20] as the boundary of a *metric compactification*. The metric compactification of a metric space (M, ρ) with the base point $x_0 \in M$ is the compactification given via Gelfand's theorem as the maximal ideal space of the C^*-algebra generated by constant functions, continuous functions vanishing at infinity, and continuous functions which form

$$\varphi_y(x) = \rho(x, x_0) - \rho(x, y)$$

for all $y \in M$. He observed that the metric compactification is naturally identified with the compactification given by M. Gromov in [6], which recently called the *horofunction compactification* (cf. §4 in [20]. See also §8.12 of Chapter II in [2]).

In [20], he also defined geodesic-like sequences in a metric space with the base point, which called *almost geodesics* (cf. §3.2). He observed that any almost geodesic admits the limit in the metric boundary. He defined *Busemann points* in the metric boundary as the limits of almost geodesics, and posed a question which asks to determine whether every point in the metric boundary of a given metric space is a Busemann point (see the paragraph after Definition 4.8 in [20]). For this problem, C. Webster and A. Winchester [23] gave geometric conditions which determine whether or not every point on the metric boundary of a graph with the standard path metric is a Busemann point, and an example of a graph which admits non-Busemann points in its metric boundary.

1.2. **Results.** Let X be a Riemann surface of type (g, n) with $2g - 2 + n > 0$. The *Teichmüller space* $T(X)$ of X is a quasiconformal deformation space of marked Riemann surfaces with same type as X. Teichmüller space $T(X)$ admits a canonical distance, called the *Teichmüller distance* d_T (cf. §2.3).

The aim of this paper is to show the following.

Theorem 1.1 (Non-Busemann points). *When $3g-3+n \geq 2$, the metric boundary of the Teichmüller space with respect to the Teichmüller distance contains non-Busemann points.*

2010 *Mathematics Subject Classification.* Primary 30F60, 32G15, Secondary 31B15, 51F99.

Key words and phrases. Teichmüller space, Teichmüller distance, extremal length, metric boundary, horofunction boundary, Busemann point.

The author is partially supported by the Ministry of Education, Science, Sports and Culture, Grant-in-Aid for Scientific Research (C), 21540177.

When $3g-3+n=1$, the Teichmüller space equipped with the Teichmüller distance is isometric to the Poincaré hyperbolic disk. Hence, every point in the metric boundary is a Busemann point. Furthermore, in this case, the metric boundary of the Teichmüller space equipped with the Teichmüller space coincides with the Thurston boundary (cf. e.g. [16]).

Recently, in [22], C. Walsh defined the horofunction boundaries for asymmetric metric spaces, and observed that the horofunction boundary of the Teichmüller space with respect to the Thurston's (non-symmetrized) Lipschitz metric is canonically identified with the Thurston boundary. He also showed that every point in the Thurston boundary is a Busemann point with respect to the Thurston's Lipschitz metric (cf. Theorem 4.1 of [22]).

The Thurston's Lipscthiz metric is the length spectrum asymmetric metric with respect to the hyperbolic lengths of simple closed curves, meanwhile the Teichmüller distance is recognized as the length spectrum metric with respect to the extremal lengths of simple closed curves via Kerckhoff's formula (cf. (2.5). See also [11]). Since hyperbolic lengths and extremal lengths are fundamental geometric quantities in the Teichmüller theory, it is natural to compare properties of these two distances. Theorem 1.1 and Walsh's results above imply that the asymptotic geometry with respect to the Teichmüller distance is more complicated than that with respect to the Thurston's Lipschitz metric.

It is known that the metric boundary of a complete CAT(0)-space consists of Busemann points (cf. Corollary II.8.20 of [2]). Therefore, we conclude the following which is already well-known (cf. [14]).

Corollary 1.1. *When $3g-3+n\geq 2$, the Teichmüller space equipped with the Teichmüller distance is not a* CAT(0)*-space.*

1.3. **The Gardiner-Masur boundary.**

Let $\mathcal{S}$ be the set of homotopy classes of non-trivial and non-peripheral simple closed curves on X. We denote by $\mathrm{Ext}_y(\alpha)$ the *extremal length* of α for $y\in T(X)$ (cf. §2.3.1). In a beautiful paper [5], F. Gardiner and H. Masur proved that the mapping

$$T(X)\ni y\mapsto [\mathcal{S}\ni\alpha\mapsto \mathrm{Ext}_y(\alpha)^{1/2}]\in \mathrm{P}\mathbb{R}_+^{\mathcal{S}}$$

is an embedding and the image is relatively compact, where $\mathbb{R}_+=\{x\geq 0\}$ and $\mathrm{P}\mathbb{R}_+^{\mathcal{S}}=(\mathbb{R}_+^{\mathcal{S}}-\{0\})/\mathbb{R}_{>0}$. The closure of the image is called the *Gardiner-Masur compactification* and the *Gardiner-Masur boundary* $\partial_{GM}T(X)$ is the complement of the image from the Gardiner-Masur compactification. They showed that the Gardiner-Masur boundary contains the space $\mathcal{PMF}$ of projective measured foliations (cf. Theorem 7.1 in [5]).

In [12], L. Liu and W. Su have shown that the horofunction boundary with respect to the Teichmüller distance is canonically identified with the Gardiner-Masur boundary of Teichmüller space. Hence, to conclude Theorem 1.1, we will show the following.

Theorem 1.2 (Non-accessibility via almost geodesics)**.** *When $3g-3+n\geq 2$, the projective class of a maximal rational measured foliation can not be the limit of any almost geodesic in the Gardiner-Masur compactification.*

In contrast, from Theorem 7.1 in [5] and Theorem 3 in [18], when a measured foliation G is either a weighted simple closed curve or a uniquely ergodic measured foliation, the projective class $[G]$ is the limit of the Teichmüller ray associated to $[G]$, and hence it is a Busemann point with respect to the Teichmüller distance.

In [17], the author have already observed that any Teichmüller geodesic ray does not converge to the projective class $[G]$ when G is a rational foliation whose support consists of at least two curves. However, the author does not know whether this induces Theorem 1.2.

This paper is organized as follows. In §2, we recall the definitions and properties of ingredients in the Teichmüller theory, including the extremal length and the Teichmüller distance. In §3, we discuss the metric boundaries of metric spaces, and check that any almost geodesic converges in the Gardiner-Masur compactification. Though this convergence follows from properties of the metric boundary and Liu and Su's work in [12], we shall give a simple proof of the convergence from the Teichmüller theory for the completeness of readers.

We treat measured foliations whose projective classes are the limits of almost geodesics in §4 and §5. Indeed, in §5, we will observe that when a measured foliation whose projective class is the limit of an almost geodesic has a foliated annulus as its component, any simple closed curve is not so *twisted* in the characteristic annulus corresponding to the foliated annulus through the almost geodesic (cf. Lemma 5.2). This is a key for getting our result. In §6, we give the proof of Theorem 1.2 by contradiction. Indeed, under the assumption that the projective class of maximal measured foliation G is the limit of an almost geodesic, we calculate the limit of a given almost geodesic, but we can check that the limit can not be equal to the boundary point induced from the intersection number function with respect to G. For getting the limit, we will apply the Kerckhoff's calculation in [9] of the extremal length along the Teichmüller ray. One of the reason why the Kerckhoff's calculation works is such *non-twisted property* of simple closed curves along the core curve of the characteristic annuli discussed in §5 (see §6.3.1).

2. EXTREMAL LENGTH AND TEICHMÜLLER THEORY

2.1. Extremal length.

Let Γ be a family of rectifiable curves on a Riemann surface R. The *extremal length* of Γ (on R) is defined by

$$\mathrm{Ext}(\Gamma)=\sup_{\rho}\frac{L_\rho(\Gamma)^2}{A(\rho)} \tag{2.1}$$

where supremum runs over all measurable conformal metric $\rho=\rho(z)|dz|^2$ and

$$L_\rho(\Gamma)=\inf_{\gamma\in\Gamma}\int_\gamma \rho(z)^{1/2}|dz| \quad \text{and} \quad A(\rho)=\int_R \rho(z)dxdy.$$

The extremal length is a *conformal invariant* in the sense that

$$\mathrm{Ext}(h(\Gamma))\le K\,\mathrm{Ext}(\Gamma) \tag{2.2}$$

for a K-quasiconformal mapping $h : R\to h(R)$, a Riemann surface R, and a family Γ of rectifiable curves on R.

Proposition 2.1 (See [1]). *Let Γ_1 and Γ_2 be two families of rectifiable curves on a Riemann surface R.*

1. *If any curve in Γ_1 is contained in a subdomain D_1 of R, the extremal length of Γ_1 on R is equal to the extremal length of Γ_1 on D_1.*
2. *If any curve in Γ_2 contains a curve in Γ_1, $\mathrm{Ext}(\Gamma_1)\le\mathrm{Ext}(\Gamma_2)$.*

(3) *If curves of* Γ_1 *and* Γ_2 *are mutually disjoint,* $\mathrm{Ext}(\Gamma_1+\Gamma_2) \geq \mathrm{Ext}(\Gamma_1) + \mathrm{Ext}(\Gamma_2)$.

where $\Gamma_1+\Gamma_2 = \{\gamma_1+\gamma_2 \mid \gamma_i \in \Gamma_i\}$.

2.1.1. *Extremal length and modulus of annulus.* For an annulus A, we denote by $\mathrm{Ext}(A)$ the extremal length of the family of simple closed curves which homotopic to the core curve of A. The *modulus* of A is the reciprocal of the extremal length of A. If A is conformally equivalent to the flat annulus $\{r_1 < |z| < r_2\}$, it holds that $\mathrm{Mod}(A) = (\log(r_2/r_1))/2\pi$.

Proposition 2.2 (cf. Proposition 9.1 of [17]). *Let A be an annulus. Let $\{\beta_k\}_{k=1}^N$ be mutually disjoint Jordan arcs joining components of ∂A such that β_{k-1} and β_{k+1} divides β_k from the other arcs (set $\beta_{N+1} = \beta_1$). Let Γ_k be the set of paths in $A - \cup_{k=1}^N \beta_k$ connecting β_k and β_{k+1}. Let ρ be the extremal metric for* $\mathrm{Ext}(A)$ *on A such that $A(\rho) = 1$. Suppose that the ρ-length of β_k is bounded for all $k = 1, \cdots, N$. Then,*

$$\mathrm{Ext}(A)^{1/2} \leq \left(\sum_{k=1}^N \mathrm{Ext}(\Gamma_k)\right)^{1/2} + B$$

where B is the totality of ρ-lengths of β_k's.

2.1.2. *Extremal lengths of simple closed curves.* For a Riemann surface Y and a simple closed curve β on Y, we define the *extremal length* $\mathrm{Ext}_Y(\beta)$ of β on Y is the extremal length of the family of rectifiable closed curves on Y homotopic to β. The extremal length is characterized geometrically as

$$\mathrm{Ext}_Y(\beta) = 1/\sup_A\{\mathrm{Mod}(A)\} = \inf_A \mathrm{Ext}(A) \tag{2.3}$$

where A runs all annuli on Y whose core is homotopic to β (cf. e.g. [9] and [21]).

2.2. **Measured foliations.** The formal product $\mathbb{R}_+ \otimes \mathcal{S} = \{t\alpha \mid t \geq 0, \alpha \in \mathcal{S}\}$ is embedded into $\mathbb{R}_+^{\mathcal{S}}$ via the intersection number function:

$$\mathbb{R}_+ \otimes \mathcal{S} \ni t\alpha \mapsto [\beta \mapsto t\, i(\alpha,\beta)] \in \mathbb{R}_+^{\mathcal{S}}.$$

The closure $\mathcal{MF} = \mathcal{MF}(X)$ of the image in $\mathbb{R}_+^{\mathcal{S}}$ is called the *space of measured foliations* on X. The *space* $\mathcal{PMF} = \mathcal{PMF}(X)$ *of projective measured foliations* is the quotient space $(\mathcal{MF} - \{0\})/\mathbb{R}_{>0}$. It is known that $\mathcal{MF}$ and $\mathcal{PMF}$ are homeomorphic to $\mathbb{R}^{6g-6+2n}$ and $S^{6g-7+2n}$, respectively (cf. [3]). It is also known that when we put $i(t\alpha, s\beta) = ts\, i(\alpha,\beta)$ for $t\alpha, s\beta \in \mathbb{R}_+ \otimes \mathcal{S}$, the intersection number function extends continuously on $\mathcal{MF} \times \mathcal{MF}$. To a measured foliation G, we associate a singular foliation and a transverse measure to the underlying foliation (cf. [3]). In this paper, we denote by $\int_\beta G$ the integration of the corresponding transverse measure over a path β.

A measured foliation G is called *rational* if G satisfies

$$i(\beta, G) = \sum_{i=1}^k w_i i(\beta, \alpha_i)$$

for some $w_i > 0$ and $\alpha_i \in \mathcal{S}$ such that $i(\alpha_i, \alpha_j) = 0$ and $\alpha_i \neq \alpha_j$ for $i, j = 1, \cdots, k$ with $i \neq j$. We write $G = \sum_{i=1}^k w_k \alpha_k$ for such measured foliation. A rational measured foliation $G = \sum_{i=1}^k w_k \alpha_k$ is *maximal* if any component of $X - \cup_{i=1}^k \alpha_i$ is a pair of pants. In this case, $k = 3g - 3 + n$.

In [9], S. Kerckhoff showed that when we put $\mathrm{Ext}_X(t\beta) = t^2\mathrm{Ext}_X(\beta)$ for $t\beta \in \mathbb{R}_+ \otimes \mathcal{S}$, the extremal length extends continuously on $\mathcal{MF}$. We define

$$\mathcal{MF}_1 = \{F \in \mathcal{MF} \mid \mathrm{Ext}_X(F) = 1\},$$

which is homeomorphic to $\mathcal{PMF}$ via the projection $\mathcal{MF} \to \mathcal{PMF}$.

In [19], Y. Minsky showed the following inequality, which recently called the *Minsky's inequality*:

$$i(F,G)^2 \leq \mathrm{Ext}_X(F)\,\mathrm{Ext}_X(G) \tag{2.4}$$

for all $F, G \in \mathcal{MF}$ (cf. Lemma 5.1 of [19]). From Theorem 5.1 in [5], Minsky's inequality is sharp in the sense that for any $G \in \mathcal{MF} - \{0\}$, there is an $F \in \mathcal{MF}$ which satisfies the equality in (2.4).

2.3. **Teichmüller space.** The *Teichmüller space* $T(X)$ of X is the set of equivalence classes of marked Riemann surfaces (Y, f) where Y is a Riemann surface and $f : X \to Y$ a quasiconformal mapping. Two marked Riemann surfaces (Y_1, f_1) and (Y_2, f_2) are *Teichmüller equivalent* if there is a conformal mapping $h : Y_1 \to Y_2$ which homotopic to $f_2 \circ f_1^{-1}$. Throughout this paper, we consider the Teichmüller space as a pointed space with the base point $x_0 = (X, id)$.

2.3.1. *Teichmüller distance and Kerckhoff's formula.* The *Teichmüller distance* between $y_1 = (Y_1, f_1)$ and $y_2 = (Y_2, f_2) \in T(X)$ is, by definition, the half of the logarithm of the extremal quasiconformal mapping between Y_1 and Y_2 preserving markings.

In [9], S. Kerckhoff gave the geometric interpretation of the Teichmüller distance by using the extremal lengths of measured foliations as follows. For $F \in \mathcal{MF}$ and $y = (Y, f) \in T(X)$, we define the extremal length of F on y by

$$\mathrm{Ext}_y(F) = \mathrm{Ext}_Y(f(F)).$$

Then, the following equality holds:

$$d_T(y_1, y_2) = \frac{1}{2}\log \sup_{H \in \mathcal{MF}-\{0\}} \frac{\mathrm{Ext}_{y_1}(H)}{\mathrm{Ext}_{y_2}(H)} = \frac{1}{2}\log \max_{H \in \mathcal{MF}_1} \frac{\mathrm{Ext}_{y_1}(H)}{\mathrm{Ext}_{y_2}(H)}. \tag{2.5}$$

The Teichmüller space is topologized with the Teichmüller distance. Under this topology, the extremal length of a measured foliation varies continuously on $T(X)$ from the conformal invariance (2.2).

2.3.2. *Quadratic differentials and Hubbard-Masur's theorem.* For a holomorphic quadratic differential $q = q(z)dz^2$ on a Riemann surface Y, we define a singular flat metric $|q| = |q(z)||dz|^2$. We call here this metric the *q-metric*.

In [7], Hubbard and Masur observed that for $y = (Y, f) \in T(X)$ and $G \in \mathcal{MF} - \{0\}$, there is a unique holomorphic quadratic differential $J_{G,y}$ on Y whose vertical foliation is equal to $f(G)$. Namely,

$$i(\beta, G) = \inf_{\beta' \in f(\beta)} \int_{\beta'} \left|\mathrm{Re}\sqrt{J_{G,y}}\right|$$

holds for all $\beta \in \mathcal{S}$. In this case, we can see that

$$\mathrm{Ext}_y(G) = \|J_{G,y}\| = \int_Y |J_{G,y}|.$$

Namely, the extremal length is the area of the $J_{G,y}$-metric. When $G = \beta \in \mathcal{S}$, we call the differential $J_{\beta,y}$ the *Jenkins-Strebel differential for* β.

2.4. **Teichmüller rays.** Let $x = (X, f) \in T(X)$ and $[G] \in \mathcal{PMF}$. By Ahlfors-Bers theorem, we can define an isometric embedding

$$[0, \infty) \ni t \mapsto R_{G,x_0}(t) \in T(X)$$

with respect to the Teichmüller distance by assigning the solution of the Beltrami equation defined by the Teichmüller Beltrami differential

$$\tanh(t)\frac{|J_{G,x_0}|}{J_{G,x_0}} \tag{2.6}$$

for $t \geq 0$. We call R_{G,x_0} the *Teichmüller (geodesic) ray associated to* $[G] \in \mathcal{PMF}$. Notice that the differential (2.6) depends only on the projective class of G. It is known that

$$\mathcal{PMF} \times [0, \infty) \ni ([G], t) \mapsto R_{G,x_0}(t) \in T(X)$$

is a homeomorphism (cf. [8]). One can see that

$$\mathrm{Ext}_{R_{G,x_0}(t)}(G) = e^{-2t}\mathrm{Ext}_{x_0}(G) \tag{2.7}$$

for $G \in \mathcal{MF}$.

2.5. **Gardiner-Masur boundary revisited.** For $y \in T(X)$, we let $K_y = e^{2d_T(x_0,y)}$. Consider a continuous function on $\mathcal{MF}$

$$\mathcal{E}_y(F) = \left(\frac{\mathrm{Ext}_y(F)}{K_y}\right)^{1/2}.$$

for $y \in T(X)$. Then, in [17], the author observed that for any $p \in \partial_{GM}T(X)$, there is a function $\mathcal{E}_p$ on $\mathcal{MF}$ such that the function $\mathcal{S} \ni \beta \mapsto \mathcal{E}_p(\beta)$ represents p and when a sequence $\{y_n\}_n \subset T(X)$ converges to p in the Gardiner-Masur compactification, there are $t_0 > 0$ and a subsequence $\{y_{n_j}\}_j$ such that $\mathcal{E}_{y_{n_j}}$ converges to $\mathcal{E}_p$ uniformly on any compact set of $\mathcal{MF}$.

3. METRIC BOUNDARY AND HOROFUNCTION BOUNDARY

3.1. **Metric boundary and horofunction boundary.** Let (M, ρ) be a locally compact metric space. Let $C(M)$ be the space of continuous functions on M, equipped with the topology of uniform convergence on compact subsets of M. Let $C_*(M)$ be the quotient space of $C(M)$ via constant functions. For $y \in M$ we set $\psi_y(x) = \rho(x, y)$. Then, $M \ni y \mapsto \psi_y$ is a continuous embedding into $C(M)$. This embedding descends a continuous embedding into $C_*(M)$. The closure $\mathcal{C\ell}(M) \subset C_*(M)$ of the image of this embedding is called the *horofunction compactification* and the complement $\mathcal{C\ell}(M) \setminus M$ is said to be the *horofunction boundary* of M (cf. [6], [2], and [20]). M. Rieffel pointed out that the metric boundary of M is canonically identified with the horofunction boundary of M as discussed in the introduction (cf. §4 in [20]).

In [12], L. Liu and W. Su showed that the horofunction compactification of the Teichmüller space with the Teichmüller distance is identified with the Gardiner-Masur compactification.

3.2. **Almost geodesics.** Let (M,ρ) be a metric space. Let $T\subset[0,\infty)$ be an unbounded set with $0\in T$. A mapping $\gamma:T\to M$ is said to be an *almost geodesic* if for any $\epsilon>0$ there is an $N>0$ such that for all $t,s\in T$ with $t\geq s\geq N$,

$$|\rho(\gamma(t),\gamma(s))+\rho(\gamma(s),\gamma(0))-t|<\epsilon$$

(cf. Definition 4.3 of [20]). By definition, any geodesic ray is an almost geodesic. When (M,ρ) is a pointed metric space, we assume in addition that $\gamma(0)$ is the base point (cf. the assumption of Lemma 4.5 in [20]). By definition, for an unbounded subset $T_0\subset T$ with $0\in T_0$, the restriction $\gamma\mid_{T_0}:T_0\to M$ is also an almost geodesic. We call the restriction a *subsequence* of an almost geodesic $\gamma:T\to M$. A point of the metric boundary or the horofunction boundary of M is said to be a *Busemann point* if it is the limit of an almost geodesic (cf. Definition 4.8 of [20]).

3.3. **Convergence of almost geodesics.** In this section, we shall check that any almost geodesic in $T(X)$ converges in the Gardiner-Masur compactification. Though this follows from a fundamental property of the metric boundary (cf. [20]) and Liu and Su's work [12], we now try to give a simple proof from the Teichmüller theory and it seems to be intriguing in itself. Notice that the author observed in [18] that any Teichmüller ray $R_{G,x}(t)$ admits the limit for all $[G]\in\mathcal{PMF}$ by the different idea.

Let $\gamma:T\to T(X)$ be an almost geodesic with the base point $x_0\in T(X)$. By definition, γ satisfies that $\gamma(0)=x_0$ and for any $\epsilon>0$, there is an N such that

$$|d_T(\gamma(t),\gamma(s))+d_T(\gamma(s),\gamma(0))-t|<\epsilon \tag{3.1}$$

for all $t\geq s\geq N$. From Kerckhoff's formula (2.5), (3.1) is equivalent to

$$e^{t-\epsilon}\leq\max_{H\in\mathcal{MF}_1}\frac{\mathrm{Ext}_{\gamma(t)}(H)^{1/2}}{\mathrm{Ext}_{\gamma(s)}(H)^{1/2}}\cdot K^{1/2}_{\gamma(s)}\leq e^{t+\epsilon}. \tag{3.2}$$

In particular, we have

$$e^{t-\epsilon}\leq K^{1/2}_{\gamma(t)}\leq e^{t+\epsilon} \tag{3.3}$$

when we set $s=t$ in (3.2). Therefore, we deduce

$$\frac{\mathrm{Ext}_{\gamma(t)}(H)^{1/2}}{\mathrm{Ext}_{\gamma(s)}(H)^{1/2}}\cdot K^{1/2}_{\gamma(s)}\leq e^{\epsilon}\cdot e^{t}\leq e^{\epsilon}\cdot e^{\epsilon}K^{1/2}_{\gamma(t)},$$

and hence

$$\mathcal{E}_{\gamma(t)}(H)\leq e^{2\epsilon}\mathcal{E}_{\gamma(s)}(H) \tag{3.4}$$

for all $H\in\mathcal{MF}$ and $t\geq s\geq N$.

We set

$$\mathcal{E}'(F)=\inf_{t\in T}\mathcal{E}_{\gamma(t)}(F)$$

for $F\in\mathcal{MF}$. From (3.4), for all $\beta\in\mathcal{S}$, the limit of any converging subsequence in $\{\mathcal{E}_{\gamma(t)}(\beta)\}_{t\in T}$ coincides with $\mathcal{E}'(\beta)$, which implies that $\gamma:T\to T(X)$ converges in the Gardiner-Masur compactification as $t\to\infty$.

4. Measured foliations as Busemann points

Let $G \in \mathcal{MF}_1$. Suppose that the projective class $[G]$ is a Busemann point in the horofunction compactification of Teichmüller space with respect to the Teichmüller metric. By definition and Liu and Su's work [12], there is an almost-geodesic $\gamma : T \to T(X)$ such that $\gamma(t) \to [G]$ in the Gardiner-Masur closure. This means that there is a $t_0 > 0$ such that $\mathcal{E}_{\gamma(t)}$ converges to $t_0\, i(\,\cdot\,, G)$ uniformly on any compact sets of $\mathcal{MF}$. We take $G_t \in \mathcal{MF}_1$ with $R_{G_t,x_0}(d_T(x_0,\gamma(t))) = \gamma(t)$.

Lemma 4.1. *Under the notation above, it holds $t_0 = 1$.*

Proof. Let G_∞ be an accumulation point of $\{G_t\}_{t\in T}$. By taking a subsequence if necessary, we may assume that G_t converges to G_∞. Let $\alpha \in \mathcal{S}$. From (2.7) and (3.3), we have

$$\begin{aligned} i(\alpha, G_\infty) &= \lim_{T\ni t\to\infty} i(\alpha, G_t) \le \lim_{T\ni t\to\infty} \mathrm{Ext}_{\gamma(t)}(\alpha)^{1/2}\mathrm{Ext}_{\gamma(t)}(G_t)^{1/2} \\ &\le e^{\epsilon} \lim_{T\ni t\to\infty} e^{-t}\mathrm{Ext}_{\gamma(t)}(\alpha)^{1/2} \le e^{2\epsilon} \lim_{T\ni t\to\infty} \mathcal{E}_{\gamma(t)}(\alpha) \\ &= e^{2\epsilon}\, t_0\, i(\alpha, G). \end{aligned} \tag{4.1}$$

Since $\epsilon > 0$ is taken arbitrary, we get

$$i(\alpha, G_\infty) \le i(\alpha, t_0 G) \tag{4.2}$$

for all $\alpha \in \mathcal{S}$. Thus, it follows from the Marden-Strebel's minimal norm property that

$$1 = \mathrm{Ext}_{x_0}(G_\infty) = \|J_{G_\infty,x_0}\| \le \|J_{t_0G,x_0}\| = t_0^2\|J_{G,x_0}\| = t_0^2, \tag{4.3}$$

and hence $t_0 \ge 1$ (see Theorem 3.2 of [13]. See also [4]).

From (3.3), by dividing every term in (3.2) by $K_{\gamma(t)}^{1/2} = e^{d_T(x_0,\gamma(t))}$ and letting $t \to \infty$, we get

$$e^{-2\epsilon} \le \max_{H\in\mathcal{MF}_1} \frac{t_0\, i(H,G)}{\mathrm{Ext}_{\gamma(s)}(H)^{1/2}} \max_{H\in\mathcal{MF}_1} \mathrm{Ext}_{\gamma(s)}(H)^{1/2} \le e^{2\epsilon} \tag{4.4}$$

for $s \ge N$. From Minsky's inequality (2.4) and Kerckhoff's formula, we have

$$\max_{H\in\mathcal{MF}_1} \frac{i(H,G)}{\mathrm{Ext}_{\gamma(s)}(H)^{1/2}} = \sup_{H\in\mathcal{MF}-\{0\}} \frac{i(H,G)}{\mathrm{Ext}_{\gamma(s)}(H)^{1/2}} = \mathrm{Ext}_{\gamma(s)}(G)^{1/2}$$

and

$$e^{s-\epsilon} \le \max_{H\in\mathcal{MF}_1} \mathrm{Ext}_{\gamma(s)}(H)^{1/2} = K(x_0,\gamma(s)) \le e^{s+\epsilon}.$$

Hence, we get

$$t_0\mathrm{Ext}_{\gamma(s)}(G)^{1/2} \le e^{3\epsilon}e^{-s}.$$

On the other hand, from the distortion property, $\mathrm{Ext}_{\gamma(s)}(G)^{1/2} \ge e^{-\epsilon}e^{-s}$ holds in general. Therefore, we have

$$t_0e^{-\epsilon}e^{-s} \le t_0\mathrm{Ext}_{\gamma(s)}(G)^{1/2} \le e^{3\epsilon}e^{-s} \tag{4.5}$$

and $t_0 \le 1$, which is what we wanted. □

From the proof of the lemma above, we also observe the following.

Corollary 4.1. *G_t converges to G as $t \to \infty$.*

Proof. Let G_∞ be an accumulation point of $\{G_t\}_{t\in T}$ as above. Recall from (4.2) and Lemma 4.1 above that $i(\alpha, G_\infty) \le i(\alpha, G)$ for all $\alpha \in \mathcal{S}$. Since $\|J_{G_\infty,x_0}\| = 1 = \|J_{G,x_0}\|$, by the calculation in (4.3) and the conclusion from the equality of the minimal norm property, we get $J_{G_\infty,x_0} = J_{G,x_0}$ and $G_\infty = G$. □

Notice from (4.5) and Lemma 4.1 that

$$\lim_{t\to\infty} \|J_{G,\gamma(t)}\| \cdot K_{\gamma(t)} = \lim_{t\to\infty} \mathrm{Ext}_{\gamma(t)}(G) \cdot K_{\gamma(t)} = 1. \tag{4.6}$$

5. Measured foliations with foliated annuli

In this section, we devote to give asymptotic behaviors of moduli of characteristic annuli corresponding to foliated annuli and the twisting number of closed geodesics on the characteristic annuli. These observations will be used for proving Theorem 1.2 in the next section.

As the previous section, we continue to suppose that the projective class $[G]$ of $G \in \mathcal{MF}_1$ is the limit of an almost geodesic $\gamma\colon T \to T(X)$. Throughout this section, we suppose in addition that G has a component of a foliated annulus with core $\alpha \in \mathcal{S}$. Namely, $G = w_0\alpha + F$ for some $w_0 > 0$ and $F \in \mathcal{MF}$. For the simplicity, we set $J_t = J_{G,\gamma(t)}$. Let $\gamma(t) = (Y_t, f_t)$ for $t \in T$ and $A_t \subset Y_t$ be the characteristic annulus of J_t for α.

We now fix a notation. For two functions $f(t)$ and $g(t)$ with variable t, $f(t) \asymp g(t)$ means that $f(t)$ and $g(t)$ are comparable in the sense that there are positive numbers B_1 and B_2 independent of the parameter t such that $B_1 g(t) \le f(t) \le B_2 g(t)$.

5.1. Moduli of characteristic annuli.

The asymptotic behavior of the modulus of A_t is given as follows.

Lemma 5.1. $\mathrm{Mod}(A_t) \asymp K_{\gamma(t)}$ *as* $t \to \infty$.

Proof. By (2.3),

$$\mathrm{Mod}(A_t) \le 1/\mathrm{Ext}_{\gamma(t)}(\alpha) \le K_{\gamma(t)}/\mathrm{Ext}_{x_0}(\alpha)$$

for all $t \in T$. In addition, by (4.6),

$$\begin{aligned} 1/\mathrm{Mod}(A_t) &= \ell_{J_t}(\alpha)/w_0 = w_0^{-2}(J_t\text{-area of } A_t) \\ &\le w_0^{-2}\|J_t\| = w_0^{-2}\mathrm{Ext}_{\gamma(t)}(G) \asymp K_{\gamma(t)}^{-1} \end{aligned}$$

as $t \to \infty$. □

5.2. Twisting numbers of paths in flat annuli.

We here define the *twisting numbers* of proper paths in flat annuli. Let $\mathbb{S}^1_L$ be the Euclidean circle of length L. Let $A = [0,m] \times \mathbb{S}^1_L$ be a flat annulus. Let $\beta \subset A$ be an (unoriented) path connecting components of ∂A. Take a universal cover $[0,m] \times \mathbb{R} \to A$. Let $\tilde\beta$ be a lift of β. Let $(0,y_1), (m,y_2) \in [0,m] \times \mathbb{R}$ be the endpoints of $\tilde\beta$. Then, we define a *twisting number* $\mathrm{tw}_A(\beta)$ of β in A by

$$\mathrm{tw}_A(\beta) = |y_1 - y_2|/L.$$

One can easily check that the twisting number is defined independently of the choice of lifts.

5.3. **Twisting numbers of geodesics.** Let $\beta \in \mathcal{S}$ with $\beta \neq \alpha$. For $t \in T$, we set β^*_t be the geodesic representative of β in Y_t with respect to the J_t-metric. If J_t admits a flat annulus whose core is homotopic to β, we choose one of closed trajectories in the flat annulus to define β^*_t.

Let $\beta^*_t \cap A_t = \{\sigma^1_s\}_{s=1}^{n_0}$ be the set of straight segment in β^*_t in the part of A_t counting multiplicity, where $n_0 = i(\beta, \alpha)$. Let $\{\sigma^2_j\}_j$ be a collection of maximal straight segments in $\beta^*_t \setminus \cup_{s=1}^{n_0} \sigma^1_s$, counting multiplicity. In this section, for a measured foliation F and a path σ transverse to the underlying foliation of F, we define $i(\sigma, F)$ as the infimum of the integrals of the transversal measure of F over all paths homotopic to σ rel endpoints.

Lemma 5.2 (Twisting number)**.** *For $s = 1, \cdots, n_0$, the twisting number of σ^1_s in A_t satisfies*

$$\mathrm{tw}_{A_t}(\sigma^1_s) = o(K_{\gamma(t)})$$

as $t \to \infty$.

Proof. When $n_0 = i(\beta, \alpha) = 0$, the geodesic representative β^*_t does not intersect the interior of A_t. Hence, the conclusion automatically holds. Therefore, we may assume that $n_0 \neq 0$.

Let $q_t = J_t / \|J_t\|$. Then, the vertical foliation V_{q_t} of q_t is equal to $\|J_t\|^{-1/2} G$ for all $t \in T$. Especially, the q_t-height w_t of the characteristic annulus A_t is equal to $\|J_t\|^{-1/2} w_0$. Let H_{q_t} be the horizontal foliation of q_t. Since each σ^1_s is a q_t-straight segment,

$$i(\sigma^j_s, V_{q_t}) = \int_{\sigma^j_s} V_{q_t} \quad \text{and} \quad i(\sigma^j_s, H_{q_t}) = \int_{\sigma^j_s} H_{q_t}$$

for $j = 1, 2$. Hence,

$$\ell_{q_t}(\beta^*_t) = \sum_{s=1}^{n_0} \sqrt{i(\sigma^1_s, H_{q_t})^2 + i(\sigma^1_s, V_{q_t})^2} + \sum_j \sqrt{i(\sigma^2_j, H_{q_t})^2 + i(\sigma^2_j, V_{q_t})^2}. \tag{5.1}$$

Since $\|q_t\| = 1$, $\ell_{q_t}(\beta^*_t) \leq \mathrm{Ext}_{\gamma(t)}(\beta)^{1/2}$ from (2.1). Therefore,

$$\begin{aligned}
\|J_t\|^{-1/2} i(\beta, G) = i(\beta, V_{q_t}) &\leq \ell_{q_t}(\beta^*_t) \\
&= \sum_{s=1}^{n_t} \sqrt{i(\sigma^1_s, H_{q_t})^2 + i(\sigma^1_s, V_{q_t})^2} + \sum_j \sqrt{i(\sigma^2_j, H_{q_t})^2 + i(\sigma^2_j, V_{q_t})^2} \\
&= \sum_{s=1}^{n_t} \sqrt{i(\sigma^1_s, H_{q_t})^2 + \|J_t\|^{-1} w_0^2} + \sum_j \sqrt{i(\sigma^2_j, H_{q_t})^2 + i(\sigma^2_j, V_{q_t})^2} \\
&\leq \mathrm{Ext}_{\gamma(t)}(\beta)^{1/2}.
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
i(\beta, G) &\leq \sum_{s=1}^{n_0} \sqrt{\|J_t\| i(\sigma_s^1, H_{q_t})^2 + w_0^2} \\
&\qquad + \sum_j \sqrt{\|J_t\| i(\sigma_j^2, H_{q_t})^2 + \|J_t\| i(\sigma_j^2, V_{q_t})^2} \\
&= \sum_{s=1}^{n_0} \sqrt{i(\sigma_s^1, H_{J_t})^2 + w_0^2} + \sum_j \sqrt{i(\sigma_j^2, H_{J_t})^2 + i(\sigma_j^2, V_{J_t})^2} \\
&\leq \|J_t\|^{1/2} \mathrm{Ext}_{\gamma(t)}(\beta)^{1/2}. \qquad (5.2)
\end{aligned}$$

From the assumption, Lemma 4.1 and (4.6),

$$\|J_t\|^{1/2} \mathrm{Ext}_{\gamma(t)}(\beta)^{1/2} = (1 + o(1)) \mathrm{Ext}_{\gamma(t)}(\beta)^{1/2} / K_{\gamma(t)}^{1/2} = (1 + o(1)) \mathcal{E}_{\gamma(t)}(\beta)$$

tends to $i(\beta, G)$ as $t \to \infty$. Since

$$i(\beta, G) = \int_{\beta_t^*} V_{J_t} = n_0 w_0 + \sum_j i(\sigma_j^2, V_{J_t}),$$

we deduce from (5.2) that the summation

$$\begin{aligned}
(5.3) \qquad & \sum_{s=1}^{n_0} \left(\sqrt{i(\sigma_s^1, H_{J_t})^2 + w_0^2} - w_0 \right) \\
&\qquad + \sum_j \left(\sqrt{i(\sigma_j^2, H_{J_t})^2 + i(\sigma_j^2, V_{J_t})^2} - i(\sigma_j^2, V_{J_t}) \right)
\end{aligned}$$

tends to zero as $t \to \infty$. Since every term in (5.3) is non-negative, we get

$$\lim_{t \to \infty} i(\sigma_s^1, H_{J_t}) = 0 \qquad (5.4)$$

for $s = 1, \cdots, n_0$.

We now fix $s = 1, \cdots, n_0$. Let $[0, w_t] \times \mathbb{R} \to [0, w_t] \times \mathbb{S}^1_{\ell_t} \cong A_t$ be the universal cover, where ℓ_t is the q_t-circumference of A_t. Let $(0, y_1)$ and (w_t, y_2) be the endpoints of a lift of σ_s^1. From the definition,

$$|y_1 - y_2| = i(\sigma_s^1, H_{q_t}).$$

Since

$$\mathrm{Mod}(A_t) = w_t / \ell_t = \|J_t\|^{-1/2} w_0 / \ell_t = (1 + o(1)) K_{\gamma(t)}^{1/2} w_0 / \ell_t,$$

from Lemma 5.1, we obtain

$$\ell_t \asymp K_{\gamma(t)}^{-1/2} \qquad (5.5)$$

for $s = 1, \cdots, n_t$. Thus, it follows from (5.4) that

$$\begin{aligned}
\mathrm{tw}_{A_t}(\sigma_s^1) &= |y_1 - y_2| / \ell_t \asymp i(\sigma_s^1, H_{q_t}) K_{\gamma(t)}^{1/2} \\
&= \|J_t\|^{-1/2} i(\sigma_s^1, H_{J_t}) K_{\gamma(t)}^{1/2} = (1 + o(1)) i(\sigma_s^1, H_{J_t}) K_{\gamma(t)} \\
&= o(K_{\gamma(t)}),
\end{aligned}$$

which implies what we wanted. □



FIGURE 1. The vertical foliation of q_t around a typical component of N_t. The component N_t^j and annuli C_i^j, $C_i^{0,j}$ in the figure are taken and used in §6.3.2.

5.4. **Twisting deformations on flat annuli.** In this section, we shall recall a canonical quasiconformal mapping of the twisting deformations along the core curve on a flat annulus (cf. [15]).

Let $A = \{e^{-2\pi m} < |z| < 1\}$ be a flat annulus of modulus m. For $t > 0$, we consider a quasiconformal self-mapping W_t of A by

$$W_t(z) = z|z|^{-\frac{i\tau}{2\pi m}}.$$

Then, the Beltrami differential of W_t is equal to

$$\frac{\overline{\partial} W_\tau}{\partial W_\tau} = \frac{-i(\tau/m)}{4\pi - i(\tau/m)} \frac{z}{\overline{z}} \frac{d\overline{z}}{dz}. \tag{5.6}$$

We can check that

$$\mathrm{tw}_A(W_\tau(\sigma)) = \mathrm{tw}_A(\sigma) - \tau. \tag{5.7}$$

Especially, when a proper path σ in A has the twist parameter τ, $\mathrm{tw}_A(W_t(\sigma)) = 0$.

6. PROOF OF THEOREM 1.2

In this section, we shall show Theorem 1.2. Throughout this section, we assume that $G = \Sigma_{i=1}^k w_i \alpha_i$ is a maximal rational foliation and $k = 3g - 3 + n \geq 2$. As before, we also assume that the projective class $[G]$ is the limit of an almost geodesic $\gamma : T \to T(X)$. We continue to use symbols given in the previous sections.

6.1. **Notation.** Let $A_{i,t} \subset Y_t$ be the characteristic annulus of $q_t = J_t/\|J_t\|$ for α_i. Let Σ_t be the critical graph of q_t and consider the $K_{\gamma(t)}^{-1/2}$-neighborhood N_t of Σ_t in Y_t with respect to the q_t-metric. Let $A_{i,t}^0 = A_{i,t} \setminus N_t$ (cf. Figure 1). Since q_t-height of $A_{i,t}$ is $(1 + o(1))K_{\gamma(t)}^{1/2} w_i$, when $t \in T$ is sufficiently large, $A_{i,t}^0$ is a well-defined foliated subannulus of $A_{i,t}$ with height

$$w''_{i,t} := (1 + o(1))K_{\gamma(t)}^{1/2} w_i - 2K_{\gamma(t)}^{-1/2} \asymp K_{\gamma(t)}^{1/2}. \tag{6.1}$$

Let $\partial A^0_{i,t}=\partial_1 A^0_{i,t}\cup\partial_2 A^0_{i,t}$. Since $\partial A^0_{i,t}$ consists of closed leaves in $A_{i,t}$ and the heights of the remaining annuli in $A_{i,t}-A^0_{i,t}$ are at most $K^{-1/2}_{\gamma(t)}$, from (5.5), the moduli of remaining annuli in $A_{i,t}-A^0_{i,t}$ are uniformly bounded, and hence

$$\mathrm{Mod}(A_{i,t})=\mathrm{Mod}(A^0_{i,t})+O(1)$$

as $t\to\infty$.

6.2. **Calculation of extremal length: Lower estimate.** We take $\beta\in\mathcal{S}$. We here devote to estimate the extremal length of β from below. From now on, we suppose that $i(\beta,G)\neq 0$.

Let $\mathcal{A}_t$ be the characteristic annulus of the Jenkins-Strebel differential $J_{\beta,\gamma(t)}$ for β. Fix $i=1,\cdots,k$. The intersection $\mathcal{A}_t\cap A^0_{i,t}$ contains at least $n_i=i(\beta,\alpha_i)$-components $\{D^i_l\}_{l=1}^{n_i}$ such that D^i_l contains a path connecting $\mathcal{A}_t\cap\partial_1 A^0_{i,t}$ and $\mathcal{A}_t\cap\partial_2 A^0_{i,t}$.

Let $\Gamma(D^i_l)$ be the family of rectifiable curves in D^i_l connecting $\mathcal{A}_t\cap\partial_1 A^0_{i,t}$ and $\mathcal{A}_t\cap\partial_2 A^0_{i,t}$. Let ρ^i_t be the restriction of the q_t-metric to $D^i=\cup_{i=1}^{n_i}D^i_l$. From (6.1), any curve in $\sum_{i=1}^{n_i}\Gamma(D^i_l)$ has ρ^i_t-length at most $n_i w''_{i,t}$. Since the critical graph of the Jenkins-Strebel differential of β on Y_t has measure zero,

$$A(\rho^i_t)=\int_{D^i}\rho^i_t\leq(|q_t|\text{-area of }A^0_{i,t})=\ell_{i,t}w''_{i,t}.$$

By the definition of the extremal length, we have

$$\begin{aligned}\mathrm{Ext}\left(\sum_{i=1}^{n_i}\Gamma(D^i_l)\right)&\geq L_{\rho^i_t}\left(\sum_{i=1}^{n_i}\Gamma(D^i_l)\right)^2/A(\rho^i_t)\geq(n_i w''_{i,t})^2/\ell_{i,t}w''_{i,t}\\&=n_i^2\mathrm{Mod}(A^0_{i,t})=n_i^2\mathrm{Mod}(A_{i,t})+O(1).\end{aligned}$$

Since any non-trivial simple closed curve in $\mathcal{A}_t$ traverses each D^i_l between $\mathcal{A}_t\cap\partial_1 A^0_{i,t}$ and $\mathcal{A}_t\cap\partial_2 A^0_{i,t}$, such simple closed curve contains a curve in $\sum_{i=1}^{n_i}\Gamma(D^i_l)$. Therefore, from (3) of Proposition 2.1, we conclude

$$\mathrm{Ext}_{\gamma(t)}(\beta)\geq\sum_{i=1}^{k}\mathrm{Ext}\left(\sum_{i=1}^{n_i}\Gamma(D^i_l)\right)\geq\sum_{i=1}^{k}n_i^2\mathrm{Mod}(A_{i,t})+O(1)\tag{6.2}$$

as $t\to\infty$.

6.3. **Calculation of extremal length: Upper estimate.** Before discussing the upper bound, we deform Y_t slightly as follows. For $i=1,\cdots,k$, we fix a component $\sigma^1_{s_i}$ of $\beta^*_t\cap A_{i,t}$. We put the Beltrami differential (5.6) on each flat annulus $A_{i,t}$ with $\tau=\mathrm{tw}_{A_{i,t}}(\sigma^1_{s_i})$. We extend the Beltrami differential to Y_t by putting 0 on the remaining part. Then, we obtain a quasiconformal deformation of Y_t with respect to the Beltrami differential to get $\gamma'(t)\in T(X)$. By Lemmas 5.1 and 5.2,

$$\mathrm{tw}_{A_{i,t}}(\sigma^1_{s_i})/\mathrm{Mod}(A_{i,t})\to 0\tag{6.3}$$

as $t\to\infty$ for all i, and hence,

$$d_T(\gamma(t),\gamma'(t))\to 0$$

when $t\to\infty$. This means that $\gamma'(t)$ has the same limit as that of $\gamma(t)$ in the Gardiner-Masur compactification. Thus, for simplifying of the notation, we may suppose that $\gamma'(t)=\gamma(t)$.

Notice from (5.7) that after this deformation, the twist parameter of each $\sigma^1_{s_i}$ is zero. Hence, any segment in $\beta^*_t \cap A_{i,t}$ has the twisting number at most one in $A_{i,t}$ for all i, because β is a simple closed curve and any two segments in $\beta^*_t \cap A_{i,t}$ do not intersect transversely in $A_{i,t}$. By taking a subsequence, we may assume that there is a (non-connected) graph Σ_0 on X such that the making $f_t : X \to Y_t$ induces an isomorphism Σ_0 and Σ_t.

6.3.1. *The idea for getting an appropriate upper bound.* To give an upper estimate, from (2.3), it suffices to construct a suitable annulus $\mathcal{A}_t$ on Y_t whose core is homotopic to $f_t(\beta)$. The procedure given here is originally due to S. Kerckhoff in [9], when a given almost geodesic γ is actually a geodesic (See also §9 of [17]). We briefly recall the case when γ is a geodesic. We first cut each characteristic annuli $A_{i,t}$ of J_t into $n_i = i(\beta, \alpha_i)$-congruent horizontal rectangles. The annulus $\mathcal{A}_t$ is made by composing appropriately such (slightly modified) n_i-congruent horizontal rectangles and ties (quadrilaterals) in N_t (cf. (6.5)). We can take such ties with uniform extremal length (cf. Claim 1). Then, by applying Proposition 2.2, we obtain an upper bound of the extremal length of $\mathcal{A}_t$.

One of the essential reason why we can get an appropriate upper bound in the case above is that, through the Teichmüller ray associated to the projective class of $G = \sum_{i=1}^k w_i\alpha_i$, there are "no" twisting deformation along α_i on the characteristic annulus, because the Teichmüller deformation is done by stretching in the horizontal and vertical directions. Indeed, the major part of the upper bound comes from the extremal length of congruent rectangles (cf. (6.8)). The 'no-twisting' property implies that the totality of the extremal lengths of such rectangles is equal to the major part of the lower estimate (6.2) (cf. (6.7)).

In the case when γ is an almost geodesic, we have already observed in (6.3) that β is not so twisted on the characteristic annuli too much. Hence, we can apply the similar argument for getting an appropriate upper bound of $\mathrm{Ext}_{y_t}(\beta)$.

6.3.2. *Ties* $\{B^j_s\}_s$. Since G is maximal, any component N^j_t $(j = 1, \cdots, 2g-2+n)$ of N_t is one of the three types: a pair of pants, an annulus with one distinguished point (a singularity of angle π or a flat point), or a half-pillow with two cone singularities of angle π (cf. Figure 2). In the the case when N^j_t is either an annulus or a half-pillow, we can deal with the same manner, and hence we now assume that N^j_t is a pair of pants.

Notice from (5.5) that the length of any component of ∂N^j_t is of order $K^{-1/2}_{\gamma(t)}$ with respect to the metric $\rho^j_t := |q_t|_{N^j_t}$. For simplifying of the notation, we assume that components of ∂N^j_t are α_{i_1}, α_{i_2}, and α_{i_3}. Then, the critical graph $\Sigma_t \cap N^j_t$ forms one of the graph in Figure 2 (cf. [3]).

We make equally spaced n_{i_l}-cuts in α_{i_l} where $n_{i_l} = i(\beta, \alpha_{i_l})$ $(l = 1, 2, 3)$. Let $C^j_{i_l}$ be a component of $N^j_t \setminus \Sigma_t$ which contains α_{i_l} in the boundary. Let $C^{0,j}_{i_l}$ be a subannulus of $C^j_{i_l}$ with height $(2K_{\gamma(t)})^{-1/2}$ and $\alpha_{i_l} \subset C^{0,j}_{i_l}$. We cut $C^{0,j}_{i_l}$ along the vertical slits with endpoints in the n_{i_l}-cuts in α_{i_l} and get a family of Euclidean rectangles. Since the circumference and the height of $C^{0,j}_{i_l}$ are of order $(K_{\gamma(t)})^{-1/2}$, the moduli of such Euclidean rectangles are uniformly bounded above and below.

Claim 1 (See Figure 3). *There is a family* $\{B^j_s\}_s$ *of (singular) quadrilaterals such that*



FIGURE 2. Graphs in N_t^j.



FIGURE 3. Typical examples of ties B_1^j and B_2^j in N_t^j. B_1^j is a 'regular' quadrilateral and B_2^j is a 'singular' quadrilateral.

(1) *$B_s^j \cap C_{i_l}^{0,j}$ is a rectangle above for all s and l,*
(2) *the arc system given by correcting cores of B_s^j's is homotopic to $\beta \cap N_t^j$, where the* core *of B_s^j is a path in B_s connecting between facing arcs in $B_s^j \cap \partial N_t^j$ and*
(3) *the extremal length of family of paths in B_s^j homotopic to the core is uniformly bounded above.*

Proof of Claim 1. Notice from (5.5) and the uniformity the moduli of $\{C_{i_l}^j\}_{l=1}^3$ that the conformal structure of N_t^j is precompact in the reduced Teichmüller space. Since the intersection numbers $\{n_{i_l}\}_{l=1}^3$ are independent of t, we can take B_s^j such that the width of each B_s^j with respect to the q_t-metric are comparable with $K_{\gamma(t)}^{-1/2}$. By definition, the $|q_t|$-area of each B_s^j is $O(K_{\gamma(t)}^{-1})$.

From the reciprocal relation between the module and the extremal length for quadrilateral or Rengel's type inequality, the extremal length $\mathrm{Ext}(B_s^j)$ of the family of paths in B_s^j homotopic to the core satisfies

$$\mathrm{Ext}(B_s^j) \leq \frac{|q_t|\text{-area}}{(|q_t|\text{-width})^2} = O(1) \tag{6.4}$$

for all s (see §4 in Chapter I of [10]). □

6.3.3. *Construction of a model $\mathcal{A}_t$ of the extremal annulus.* We divide each $A_{i,t}$ into congruence $n_i = i(\beta, \alpha_i)$-rectangles $\{R_{i,l}\}_{l=1}^{n_i}$ via proper horizontal segments. We may assume that for any l and j, there is an s such that $R_{i,l} \cap C_i^{0,j}$ is congruent



FIGURE 4. Rectangles $R_{i,l}$ and $R_{i,l}^0$ in the annulus $A_{i,t}$.

to $B_s^j \cap C_i^{0,j}$. We set

$$R_{i,l}^0 = R_{i,l} \setminus (N_t \setminus \cup_j C_i^{0,j})$$

(cf. Figure 4). Since twisting numbers of segments in $\beta_t^* \cap A_{i,t}$ on each $A_{i,t}$ are at most one for all i, from (5.6) and the Dehn-Thurston's parametrization of simple closed curves (cf. [3]), we can glue all $A_{i,t}$ and N_t appropriately at the part $C_i^{0,j}$ to get a Riemann surface Y_t' and an annulus

$$\mathcal{A}_t' = (\cup_{i,l} R_{i,l}^0) \cup (\cup_{j,s} B_s^j) \tag{6.5}$$

such that after deforming Y_t' by a quasiconformal mapping with maximal dilatation $1 + o(1)$, we obtain Y_t and the core of the image $\mathcal{A}_t$ of the annulus $\mathcal{A}_t'$ is homotopic to $f_t(\beta)$. Thus, we conclude

$$\mathrm{Ext}_{y_t}(\beta) = \mathrm{Ext}_{Y_t}(f_t(\beta)) \le \mathrm{Ext}(\mathcal{A}_t) = (1 + o(1))\mathrm{Ext}(\mathcal{A}_t') \tag{6.6}$$

as $t \to \infty$. Therefore, to get the upper estimate of the extremal length of β on y_t, it suffices to give an upper estimate of the extremal length of $\mathcal{A}_t'$.

6.3.4. *Estimate of extremal length of* $\mathcal{A}_t'$. Let $\rho_t^{\mathcal{A}}$ be the extremal metric on $\mathcal{A}_t'$ for the extremal length $\mathrm{Ext}(\mathcal{A}_t')$ with $A(\rho_t^{\mathcal{A}}) = 1$. Let $\{\mathcal{S}_u\}_u$ be a collection of all rectangles of form $C_i^{0,j} \cap B_s^j$ for all i, j, s. By the same argument as Claim 1 in §9.6, we can see the following.

Claim 2. *For any u, there is a vertical line η_u' in $\mathcal{S}_u$ such that*

$$\sum_u \ell_{\rho_t^{\mathcal{A}}}(\eta_u') = O(1)$$

as $t \to \infty$.

Let us continue the calculation. Let $\{D_u\}_u$ be a collection of components of $\mathcal{A}_t' \setminus \cup_u \eta_u'$. By labeling correctly, ∂D_u contains η_u' and η_{u+1}', where η_u' is labeled cyclically in u. By definition, each D_u is contained in either $R_{i,l}$ or B_s^j for some i, l, j, s. Let $\Gamma(D_u)$ be the family of paths connecting vertical segments η_u' and η_{u+1}'. Let $\Gamma(R_{i,l})$ be the family of paths in $R_{i,l}$ connecting vertical boundary segments. Since

$$\mathrm{Ext}(\Gamma(R_{i,l})) = (\text{height of } A_{i,t})/(\ell_{i,t}/n_i) = n_i \mathrm{Mod}(A_{i,t}), \tag{6.7}$$

by (1) and (2) of Proposition 2.2 and Claim 2, we have

$$
\begin{aligned}
\operatorname{Ext}(\mathcal{A}'_t)^{1/2} &= \left(\sum_u \operatorname{Ext}(\Gamma(D_u))\right)^{1/2} + O(1) \\
&\leq \left(\sum_{D_u \subset R_{i,l}} \operatorname{Ext}(\Gamma(R_{i,l})) + \sum_{D_u \subset B_s^j} \operatorname{Ext}(B_s^j)\right)^{1/2} + O(1) \\
&\leq \left(\sum_{D_u \subset R_{i,l}} n_i \operatorname{Mod}(A_{i,t}) + O(1)\right)^{1/2} + O(1) \\
&= \left(\sum_{i=1}^k n_i^2 \operatorname{Mod}(A_{i,t}) + O(1)\right)^{1/2} + O(1). \qquad (6.8)
\end{aligned}
$$

Thus we get an upper bound of the extremal length of $\mathcal{A}'_t$ as we desired.

6.4. **Conclusion.** From Lemma 5.1, by taking a subsequence if necessary, we may assume that $\operatorname{Mod}(A_{i,t})/K_{\gamma(t)}$ tends to a positive number M_i for any $i = 1, \cdots, k$. From (6.2), (6.6), and (6.8), we deduce that

$$
i(\beta, G) = \lim_{t\to\infty} \left(\frac{\operatorname{Ext}_{\gamma(t)}(\beta)}{K_{\gamma(t)}}\right)^{1/2} = \left(\sum_{i=1}^k n_i^2 M_i\right)^{1/2} = \left(\sum_{i=1}^k M_i i(\beta, \alpha_i)^2\right)^{1/2} \tag{6.9}
$$

for all $\beta \in \mathcal{S}$. From the density of $\mathbb{R}_+ \otimes \mathcal{S}$ in $\mathcal{MF}$, the above also holds for all measured foliations. Thus, for $x, y > 0$ and $\beta_1, \beta_2 \in \mathcal{S}$ with $i(\beta_1, \beta_2) = 0$, by substituting $\beta = x\beta_1 + y\beta_2$ to (6.9), we get

$$
\begin{aligned}
\left(\sum_{i=1}^k M_i n_{1,i}^2\right) x^2 + 2\left(\sum_{i=1}^k M_i n_{1,i} n_{2,i}\right) xy + \left(\sum_{i=1}^k M_i n_{2,i}^2\right) y^2 \\
= (x i(\beta_1, G) + y i(\beta_2, G))^2,
\end{aligned}
$$

where $n_{j,i} = i(\beta_j, \alpha_i)$. Therefore, the discriminant of the quadratic form above is zero. Namely, we have

$$
\left(\sum_{i=1}^k M_i n_{1,i} n_{2,i}\right)^2 - \left(\sum_{i=1}^k M_i n_{1,i}^2\right)\left(\sum_{i=1}^k M_i n_{2,i}^2\right) = 0,
$$

for all such $\beta_1, \beta_2 \in \mathcal{S}$. Hence, two vectors

$$
(\sqrt{M_1} n_{1,1}, \cdots, \sqrt{M_k} n_{1,k}), \quad (\sqrt{M_1} n_{2,1}, \cdots, \sqrt{M_k} n_{2,k}),
$$

are parallel for all $\beta_1, \beta_2 \in \mathcal{S}$ with $i(\beta_1, \beta_2) = 0$. However, this is impossible as we already observed in §6 of [17].

References

[1] L. V. Ahlfors, *Lectures on Quasiconformal Mappings*, Van Nostrand Mathematical Studies, vol. **10**. D. Van Nostrand Co., Inc., Toronto, Ont.-New York-London (1966) .

[2] M. Bridson and A. Haefliger, *Metric spaces of Non-positive curvature*, Grundlehren der mathematischen Wissenschaften **319**, Springer Verlag (1999).

[3] A. Douady, A. Fathi, D. Fried, F. Laudenbach, V. Poénaru, and M. Shub, *Travaux de Thurston sur les surfaces*, Séminaire Orsay (seconde édition). Astérisque No. 66-67, Société Mathématique de France, Paris (1991).

[4] F. Gardiner, Measured foliations and the minimal norm property for quadratic differentials, Acta Math. **152** (1984), no. 1-2, 57–76.

[5] F. Gardiner and H. Masur, Extremal length geometry of Teichmüller space. Complex Variables Theory Appl. **16** (1991), no. 2-3, 209–237.

[6] M. Gromov, Hyperbolic manifolds, groups and actions, In *Riemann surfaces and related topics*, Proceedings of the 1978 Stony Brook Conference, 182–213, Princeton University Press (1981).

[7] J. Hubbard, and H. Masur, Quadratic differentials and foliations, Acta Math. **142** (1979), no. 3-4, 221–274.

[8] Y. Imayoshi and M. Taniguchi, *Introduction to Teichmüller spaces*, Springer-Verlag (1992).

[9] S. Kerckhoff, The asymptotic geometry of Teichmüller space, Topology **19** (1980), 23–41.

[10] O. Lehto and K. Virtanen, *Quasiconformal mappings in the plane, Second edition*, Die Grundlehren der mathematischen Wissenschaften, Band **126**. Springer-Verlag, New York-Heidelberg, (1973).

[11] L. Liu, A. Papadopoulos, W. Su and G. Théret, Length spectra and the Teichmüller for surfaces with boundary, Monatsh Math **161**, 295–311 (2010).

[12] L. Liu and W. Su, The horofunction compactification of Teichmüller metric, preprint, ArXiv.org : `http://arxiv.org/abs/1012.0409`.

[13] A. Marden, and K. Strebel, The height theorem for quadratic differentials on Riemann surfaces, Acta. Math **153** (1984), 153–211.

[14] H. Masur, On a class of geodesics in Teichmüller space. Ann. of Math. **102** (1975), 205–221.

[15] A. Marden and H. Masur, A foliation of Teichmüller space by twist invariant disks, Math. Scand. **36** (1975), 211–228.

[16] H. Miyachi, On the Gardiner-Masur boundary of Teichmüller spaces Proceedings of the 15th ICFIDCAA Osaka 2007, OCAMI Studies 2 (2008), 295–300.

[17] ———, Teichmüller rays and the Gardiner-Masur boundary of Teichmüller space. Geom. Dedicata **137** (2008), 113–141.

[18] ———, Teichmüller rays and the Gardiner-Masur boundary of Teichmüller space II, submitted.

[19] Y. Minsky, Teichmüller geodesics and ends of hyperbolic 3-manifolds. Topology **32**, 625–647 (1993).

[20] M. Rieffel, Group C^*-algebra as compact quantum metric spaces, Doc. Math. 7 (2002), 605–651.

[21] K. Strebel, *Quadratic Differentials*, Springer Verlag, Berlin and New York (1984).

[22] C. Walsh, The horoboundary and isometry group of Thurson's Lipschitz metric, preprint, ArXiv.org : `http://arxiv.org/abs/1006.2158`

[23] C. Webster and A. Winchester, Busemann points of infinite graphs, Trans. Amer. Math. Soc. **358** (2006), 4209–4224.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, MACHIKANEYAMA 1-1, TOYONAKA, OSAKA 560-0043, JAPAN